MUST USE GUIDE

This chart MUST be used when referencing any questions related to pain scales. Do NOT fill out this for. It is for reference purposes only.

Func	ctional Pain Scale	10	Worst imaginable pain . Causes you to be completely incapacitated and barely able to talk. Requires
10	Worst imaginable pain		immediate emergency hospitalization.
9		8-9	Pain that causes disability between levels 7 and 10. Nearing need for hospitalization.
8 7	Severely disabling pain	7	Severely disabling pain . You cannot use or move the painful area. You have difficulty talking and concentrating on anything but the pain. Needing to lie down and/or
6			pain-related tearfulness are common at this level of pain.
5	Very disabling pain	6	Pain that causes disability between levels 5 and 7.
4		5	Very disabling pain . Causes great difficulty moving or applying any strength through the painful area. You are unable to complete the
3	Functionally disabling pain		current activity.
2		4	Pain that causes disability between levels 3 and 5
1		3	Functionally disabling pain . Pain that is starting to affect your ability to perform the current activity. (i.e., decreased movement, decreased
0.25	Non-disabling pain		speed, and/or the need to briefly rest and/or stretch in order to continue completing the current activity)
U	No pain	2.75 To 0.25	Non disabling pain. The pain is present, but not yet at a level which limits you
	MATHESON	0.25	from performing the current activity. No pain or discomfort.
Sec	www.roymatheson.com 800-443-7690 or 1-603-358-6525 Copyright © 2003 by ional Outcomes Rehabilitation Services, Inc.		

Patient Summany Form

Ins	tru	CTL	nn	S

Patient Summary Form PSF-750 (Rev:2/18	/2009)				timeli	ne and fax to	this form within the specified o the specified fax number
Patient Information			ale			licated on P n previously	lan Summary or plan infor- provided.
Patient name Last First			, l	Patient date of birth	*Fax	number may	vary by plan.
				Futient dute of binn		1	
Patient address		City				State	Zip code
Patient insurance ID#	Health plan			Group nu	ımber		
Referring physician (if applicable)	Date referral is:	sued (if applicabl	e)	Referral	number (if applica	able)	
Provider Information							
4 Name of the billing provider or facility (- 1 - 11)				2 Federal for ID/TIM) of a	atitu in hav #4		
1. Name of the billing provider or facility (as it will appear on the claim			_	2. Federal tax ID(TIN) of e	-		
		2 DC 3 P	T 4 OT	5 Both PT and OT 6	Home Care 7	ATC 8	MT 9 Other
3. Name and credentials of the individual performing the service(5)						
4 Alternate name (if any) of ontity in her #1			h				Dhana number
4. Alternate name (if any) of entity in box #1		5. NPI of entity in	box #1			1 6	. Phone number
7. Address of the billing provider or facility indicated in box #1			8. City			9. State	10. Zip code
Provider Completes This Section:				Date of Surgery			nosis (ICD code) se ensure all digits are
Date you want <i>THIS</i> submission to begin: Cause o	f Current Episo	ode	(entered accurately
(1) Traumation	\cap			/pe of Surgery	1°		
(2) Unspecifi	×			CL Reconstruction	2°		
Patient Type (3) Repetitive	X	vehicle	(2) R	otator Cuff/Labral Repa			•
1 New to your office	0		<u>(3)</u> т	endon Repair	3°		
Est'd, new injury			(4) s	pinal Fusion	5		•
3 Est'd, new episode			б ј	pint Replacement	4°		
(4) Est'd, continuing care			6 c	ther			•
Nature of Condition	DC C	ONLY		C	rent Function	al Maaai	
(1) Initial onset (within last 3 months)	Anticipated	CMT Level					
(2) Recurrent (multiple episodes of < 3 months)	98940	() 98942		Neck Index	DAS	н 厂	(other)
(3) Chronic (continuous duration > 3 months)	98941	98943		Back Index	LEF	s	(other)
Patient Completes This Section:	ms began on	:		In	dicate where y	ou have	pain or other symptom
(Please fill in selections completely)	augun en				5.7		(20)
1. Briefly describe your symptoms:					All	>	(The p
					1 X	XI	LAXIL
2. How did your symptoms start?					1754	161	114:411
					They I	aw law	Tel () for
3. Average pain intensity:	~ ~ ~	~ ~ ~	~		Julia	(Y. A.J
Last 24 hours: no pain $\begin{pmatrix} 0 \\ 1 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix}$	4 5 6 (789	10	worst pain			(1)(1)
Past week: no pain $\begin{pmatrix} 0 \\ 1 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix}$	(4) (5) (6) (7) (8) (9)	(10)	worst pain)/X<() ((
4. How often do you experience your symp		\cap			and Gent	-	See Car
(1) Constantly (76%-100% of the time) (2) Frequent	y (51%-75% of the	e time) (3) C	ccasiona)	ly (26% - 50% of the tin	ne) (4) Interm	ittently (0%	6-25% of the time)
5. How much have your symptoms interfer		/			ork outside the he	ome and h	ousework)
$\begin{pmatrix} 1 \end{pmatrix}$ Not at all $\begin{pmatrix} 2 \end{pmatrix}$ A little bit $\begin{pmatrix} 3 \end{pmatrix}$ Mode	rately (4) Q	uite a bit	5) Extre	mely			
6. How is your condition changing, since o	are began at	this facility	?				<u>^</u>
0 N/A — This is the initial visit 1 Much	worse (2) Wors	e (3) A little	worse (4) No change (5) A	little better	Better	(7) Much better
7. In general, would you say your overall h	ealth right n	ow is		-			
(1) Excellent (2) Very good (3) Good	\sim		5) Poor				
0 0 0	\cup	(Data		
Patient Signature: X					Date:		

Back Index

Form BI100

Patient Name

rev 3/27/2003

Date

This questionnaire will give your provider information about how your back condition affects your everyday life. Please answer every section by marking the one statement that applies to you. If two or more statements in one section apply, please mark the one statement that most closely describes your problem.

Pain Intensity

- ① The pain comes and goes and is very mild.
- ① The pain is mild and does not vary much.
- ② The pain comes and goes and is moderate.
- ③ The pain is moderate and does not vary much.
- The pain comes and goes and is very severe.
- ⑤ The pain is very severe and does not vary much.

Sleeping

- I get no pain in bed.
- ① I get pain in bed but it does not prevent me from sleeping well.
- 2 Because of pain my normal sleep is reduced by less than 25%.
- 3 Because of pain my normal sleep is reduced by less than 50%.
- ④ Because of pain my normal sleep is reduced by less than 75%.
- ⑤ Pain prevents me from sleeping at all.

Sitting

- ① I can sit in any chair as long as I like.
- ① I can only sit in my favorite chair as long as I like.
- ② Pain prevents me from sitting more than 1 hour.
- ③ Pain prevents me from sitting more than 1/2 hour.
- ④ Pain prevents me from sitting more than 10 minutes.
- ⑤ I avoid sitting because it increases pain immediately.

Standing

- I can stand as long as I want without pain.
- ① I have some pain while standing but it does not increase with time.
- ② I cannot stand for longer than 1 hour without increasing pain.
- ③ I cannot stand for longer than 1/2 hour without increasing pain.
- (4) I cannot stand for longer than 10 minutes without increasing pain.
- ⑤ I avoid standing because it increases pain immediately.

Walking

- I have no pain while walking.
- ① I have some pain while walking but it doesn't increase with distance.
- ② I cannot walk more than 1 mile without increasing pain.
- ③ I cannot walk more than 1/2 mile without increasing pain.
- I cannot walk more than 1/4 mile without increasing pain.
- ⑤ I cannot walk at all without increasing pain.

Personal Care

- ① I do not have to change my way of washing or dressing in order to avoid pain.
- ① I do not normally change my way of washing or dressing even though it causes some pain.
- ② Washing and dressing increases the pain but I manage not to change my way of doing it.
- ③ Washing and dressing increases the pain and I find it necessary to change my way of doing it.
- ④ Because of the pain I am unable to do some washing and dressing without help.
- (5) Because of the pain I am unable to do any washing and dressing without help.

Lifting

- I can lift heavy weights without extra pain.
- ① I can lift heavy weights but it causes extra pain.
- ② Pain prevents me from lifting heavy weights off the floor.
- ③ Pain prevents me from lifting heavy weights off the floor, but I can manage if they are conveniently positioned (e.g., on a table).
- ④ Pain prevents me from lifting heavy weights off the floor, but I can manage light to medium weights if they are conveniently positioned.
- ⑤ I can only lift very light weights.

Traveling

- I get no pain while traveling.
- ① I get some pain while traveling but none of my usual forms of travel make it worse.
- 2 I get extra pain while traveling but it does not cause me to seek alternate forms of travel.
- ③ I get extra pain while traveling which causes me to seek alternate forms of travel.
- ④ Pain restricts all forms of travel except that done while lying down.
- ⑤ Pain restricts all forms of travel.

Social Life

- My social life is normal and gives me no extra pain.
- ① My social life is normal but increases the degree of pain.
- ② Pain has no significant affect on my social life apart from limiting my more energetic interests (e.g., dancing, etc).
- ③ Pain has restricted my social life and I do not go out very often.
- ④ Pain has restricted my social life to my home.
- ⑤ I have hardly any social life because of the pain.

Changing degree of pain

- O My pain is rapidly getting better.
- ① My pain fluctuates but overall is definitely getting better.
- ② My pain seems to be getting better but improvement is slow.
- ③ My pain is neither getting better or worse.
- My pain is gradually worsening.
- S My pain is rapidly worsening.



Neck Index

Form N1-100

Patient Name

rev 3/27/2003

Date

This questionnaire will give your provider information about how your neck condition affects your everyday life. Please answer every section by marking the one statement that applies to you. If two or more statements in one section apply, please mark the one statement that most closely describes your problem.

Pain Intensity

- I have no pain at the moment.
- ① The pain is very mild at the moment.
- ② The pain comes and goes and is moderate.
- The pain is fairly severe at the moment.
- ④ The pain is very severe at the moment.
- (5) The pain is the worst imaginable at the moment.

Sleeping

- I have no trouble sleeping.
- ① My sleep is slightly disturbed (less than 1 hour sleepless).
- ② My sleep is mildly disturbed (1-2 hours sleepless).
- ③ My sleep is moderately disturbed (2-3 hours sleepless).
- ④ My sleep is greatly disturbed (3-5 hours sleepless).
- (5) My sleep is completely disturbed (5-7 hours sleepless).

Reading

- I can read as much as I want with no neck pain.
- ① I can read as much as I want with slight neck pain.
- ② I can read as much as I want with moderate neck pain.
- ③ I cannot read as much as I want because of moderate neck pain.
- ④ I can hardly read at all because of severe neck pain.
- ⑤ I cannot read at all because of neck pain.

Concentration

- ① I can concentrate fully when I want with no difficulty.
- ① I can concentrate fully when I want with slight difficulty.
- ② I have a fair degree of difficulty concentrating when I want.
- ③ I have a lot of difficulty concentrating when I want.
- (4) I have a great deal of difficulty concentrating when I want.
- I cannot concentrate at all.

Work

- I can do as much work as I want.
- ① I can only do my usual work but no more.
- $\ensuremath{\textcircled{O}}$ I can only do most of my usual work but no more.
- ③ I cannot do my usual work.
- ④ I can hardly do any work at all.
- ⑤ I cannot do any work at all.

Personal Care

- O I can look after myself normally without causing extra pain.
- ① I can look after myself normally but it causes extra pain.
- ② It is painful to look after myself and I am slow and careful.
- ③ I need some help but I manage most of my personal care.
- ④ I need help every day in most aspects of self care.
- (5) I do not get dressed, I wash with difficulty and stay in bed.

Lifting

- I can lift heavy weights without extra pain.
- ① I can lift heavy weights but it causes extra pain.
- Pain prevents me from lifting heavy weights off the floor, but I can manage if they are conveniently positioned (e.g., on a table).
- ③ Pain prevents me from lifting heavy weights off the floor, but I can manage light to medium weights if they are conveniently positioned.
- I can only lift very light weights.
- (5) I cannot lift or carry anything at all.

Driving

- I can drive my car without any neck pain.
- ① I can drive my car as long as I want with slight neck pain.
- ② I can drive my car as long as I want with moderate neck pain.
- ③ I cannot drive my car as long as I want because of moderate neck pain.
- ④ I can hardly drive at all because of severe neck pain.
- ⑤ I cannot drive my car at all because of neck pain.

Recreation

- O I am able to engage in all my recreation activities without neck pain.
- ① I am able to engage in all my usual recreation activities with some neck pain.
- ② I am able to engage in most but not all my usual recreation activities because of neck pain.
- 3 I am only able to engage in a few of my usual recreation activities because of neck pain.
- I can hardly do any recreation activities because of neck pain.
- ⑤ I cannot do any recreation activities at all.

Headaches

- I have no headaches at all.
- ① I have slight headaches which come infrequently.
- ② I have moderate headaches which come infrequently.
- ③ I have moderate headaches which come frequently.
- I have severe headaches which come frequently.
- ⑤ I have headaches almost all the time.







The STarT Back Musculoskeletal Screening Tool

Patient name: Date:

Thinking about the last 2 weeks tick your response to the following questions:

		Disagree	Agree
1	My pain has spread at some time in the past 2 weeks		
2	In addition to my main pain, I have had pain elsewhere in the last 2 weeks		
3	In the last 2 weeks, I have only walked short distances because of my pain		
4	In the last 2 weeks, I have dressed more slowly than usual because of my pain		
5	It's really not safe for a person with a condition like mine to be physically active		
6	Worrying thoughts have been going through my mind a lot of the time in the last 2 weeks		
7	I feel that my pain is terrible and that it's never going to get any better		
8	In general in the last 2 weeks, I have not enjoyed all the things I used to enjoy		

9. Overall, how bothersome has your pain been in the last 2 weeks?

Not at all	Slightly	Moderately	Very much	Extremely
0	0 0	0		

Originally developed by: © Keele University 01/08/07 Funded by Arthritis Research UK